


QL
401
V4X
MOLL
IE

VELIGER



A Quarterly published by
CALIFORNIA MALACOOLOGICAL SOCIETY, INC.
Berkeley, California
R. Stohler, Founding Editor

Volume 33 July 2, 1990 Number 3

CONTENTS

Photic vesicles. RICHARD M. EAKIN	209
Continuous reproduction and episodic recruitment of <i>Lacuna vincta</i> (Montagu, 1803) in the Gulf of Maine. EDWARD J. MANEY JR. AND JOHN P. EBERSOLE	215
Recruitment of the estuarine soft-bottom bivalve <i>Polymesoda caroliniana</i> and its influence on the vertical distribution of adults. DAN C. MARELLI	222
Additional opisthobranch mollusks from Oregon, with a review of deep-water records and observations on the fauna of the south coast. JEFFREY H. R. GODDARD	230
Temporary northern range extension of the squid <i>Loligo opalescens</i> in Southeast Alaska. BRUCE L. WING AND ROGER W. MERCER	238
A review of the Recent eastern Pacific Acanthochitoninae (Mollusca: Polyplacophora: Cryptoplacidae) with the description of a new genus, <i>Americhiton</i> . G. THOMAS WATTERS	241
New species of Late Cretaceous Cypraeacea (Mollusca: Gastropoda) from California and Mississippi, and a review of Cretaceous cypraeaceans of North America. LINDSEY T. GROVES	272

CONTENTS — *Continued*

The Veliger (ISSN 0042-3211) is published quarterly on the first day of January, April, July, and October. Rates for Volume 33 are \$28.00 for affiliate members (including domestic mailing charges) and \$56.00 for libraries and nonmembers (including domestic mailing charges). For subscriptions sent to Canada and Mexico, add US \$4.00; for subscriptions sent to addresses outside of North America, add US \$8.00, which includes air-expedited delivery. Further membership and subscription information appears on the inside cover. The Veliger is published by the California Malacozoological Society, Inc., % Museum of Paleontology, University of California, Berkeley, CA 94720. Second Class postage paid at Berkeley, CA and additional mailing offices. POSTMASTER: Send address changes to The Veliger, Museum of Paleontology, University of California, Berkeley, CA 94720.



Scope of the journal

The Veliger is open to original papers pertaining to any problem concerned with mollusks.

This is meant to make facilities available for publication of original articles from a wide field of endeavor. Papers dealing with anatomical, cytological, distributional, ecological, histological, morphological, physiological, taxonomic, evolutionary, etc., aspects of marine, freshwater, or terrestrial mollusks from any region will be considered. Short articles containing descriptions of new species or lesser taxa will be given preferential treatment in the speed of publication provided that arrangements have been made by the author for depositing the holotype with a recognized public Museum. Museum numbers of the type specimen must be included in the manuscript. Type localities must be defined as accurately as possible, with geographical longitudes and latitudes added.

Very short papers, generally not exceeding 500 words, will be published in a column entitled "NOTES, INFORMATION & NEWS"; in this column will also appear notices of meetings, as well as news items that are deemed of interest to our subscribers in general.

Editor-in-Chief

David W. Phillips, 2410 Oakenshield Road, Davis, CA 95616, USA

Editorial Board

Hans Bertsch, National University, Inglewood, California
James T. Carlton, University of Oregon
Eugene V. Coan, Research Associate, California Academy of Sciences, San Francisco
J. Wyatt Durham, University of California, Berkeley
William K. Emerson, American Museum of Natural History, New York
Terrence M. Gosliner, California Academy of Sciences, San Francisco
Cadet Hand, University of California, Berkeley
Carole S. Hickman, University of California, Berkeley
David R. Lindberg, University of California, Berkeley
James H. McLean, Los Angeles County Museum of Natural History
Frank A. Pitelka, University of California, Berkeley
Peter U. Rodda, California Academy of Sciences, San Francisco
Clyde F. E. Roper, National Museum of Natural History, Washington
Barry Roth, Santa Barbara Museum of Natural History
Judith Terry Smith, Stanford University
Ralph I. Smith, University of California, Berkeley
Wayne P. Sousa, University of California, Berkeley
T. E. Thompson, University of Bristol, England

in memoriam

Membership and Subscription

Affiliate membership in the California Malacozoological Society is open to persons (no institutional memberships) interested in any aspect of malacology. As an affiliate member, a person may subscribe to The Veliger for US \$28.00 (Volume 33), which now *includes* mailing charges to domestic addresses. There is a one-time membership fee of US \$2.00, after payment of which, membership is maintained in good standing by the timely renewal of the subscription; a reinstatement fee of US \$3.00 will be required if membership renewals do not reach the Society on or before October 1 preceding the start of the new Volume. If a receipt is required, a self-addressed, stamped envelope (or in the case of foreign members, the envelope and two International Postal Reply coupons) should be included with the membership or subscription request.

The annual subscription rate to The Veliger for libraries and nonmembers is US \$56.00 (Volume 33), which now *includes* mailing charges to domestic addresses.

For subscriptions sent to Canada and Mexico, add US \$4.00; for subscriptions sent to addresses outside of North America, add US \$8.00, which includes air-expedited delivery.

Memberships and subscriptions are by Volume only (January 1 to October 1) and are payable in advance to California Malacozoological Society, Inc. Single copies of an issue are US \$25.00 plus postage.

Send all business correspondence, including subscription orders, membership applications, payments for them, changes of address, to: The Veliger, Museum of Paleontology, 3 Earth Sciences Bldg., University of California, Berkeley, CA 94720, USA.

Send manuscripts, proofs, books for review, and correspondence regarding editorial matters to: Dr. David W. Phillips, Editor, 2410 Oakenshield Road, Davis, CA 95616, USA.

Photic Vesicles¹

by

RICHARD M. EAKIN

Department of Molecular and Cell Biology, University of California,
Berkeley, California 94720, USA

Abstract. A summary is presented of various investigations on unique cytoplasmic organelles—termed photic vesicles—in the eyes of gastropod mollusks. Emphasis is given to the research of the author and his associates and to that to Tomiyuki Hara and his colleagues at Osaka University. Some unpublished findings of the Osaka workers are included. The following aspects of the vesicles are covered: structure, origin, fate, and function as transporters of the photopigment retinochrome and calcium. The data are synthesized into a scenario, with a heretofore unpublished diagram, on the cycling of retinochrome in which photic vesicles play a strategic role.

Photic vesicles are remarkable cytoplasmic organelles in the eyes of gastropod mollusks. They are unique in aggregating in very large numbers in the cell bodies of type I photosensory cells (Figures 1, 3), and in carrying a photopigment, hence the term “photic vesicles.” They have been studied mostly in pulmonates (*e.g.*, snails and slugs—referenced below), but they occur also in opisthobranchs (*e.g.*, nudibranchs—EAKIN *et al.*, 1967), and in prosobranchs (*e.g.*, periwinkles—MAYES & HERMANS, 1973).

Early History

The vesicles were first observed by two groups of European investigators studying the ultrastructure of the eyes of a snail, *Helix pomatia*: RÖHLICH & TÖRÖK (1963), who described how the vesicles—called *Elementarkörperchen*—form a crystalline mass (*Biokristall*) and SCHWALBACH *et al.* (1963) who regarded the vesicles as a foam (*Schaumstruktur*). The truly vesicular nature of the organelle was confirmed and further described in another snail, *Helix aspersa*, by Mrs. Jean Brandenburger, my research associate for many years, and me (EAKIN & BRANDENBURGER, 1967a).

Recently I learned in a personal communication from Professor Tomiyuki Hara of Osaka University that he and two colleagues, Y. Koshida and A. Tanaka, had studied the eyes of a land snail, *Euhadra callizona amaliae*, in 1963 and observed masses of vesicles in the cell bodies of the photosensory cells. Their findings on the vesicles, however, were not published. I have now received from him an

unpublished manuscript with six plates of electron micrographs. It contains much information on the structural and biochemical features of a snail’s eye that was regrettably not entered into the scientific literature. The authors had measured the vesicles (40–50 nm in diameter) and had shown the beautiful paracrystalline arrangement of them when compacted.

Dr. Jane Westfall and I had observed the vesicles in *Helix aspersa* in a survey of invertebrate photoreceptors that led to a 1962 symposium paper (EAKIN, 1963), but it was not until 1967 that a description of them was published (EAKIN & BRANDENBURGER, 1967a). In that year we suggested the possibility that the vesicles may carry a photopigment (EAKIN & BRANDENBURGER, 1967b). At the same time a similar function was postulated by HARA *et al.* (1967) in an excellent paper on photopigments in the retina of an octopus. In the discussion they analogized the gastropod vesicles with lamellated bodies in cephalopod photosensory cells shown by them to contain a new pigment termed retinochrome. We gave to the vesicular organelle the name photic vesicle (EAKIN & BRANDENBURGER, 1970).

Structure

The vesicles are spherical when not compressed, and uniform in size (diameter 80–85 nm in *Helix aspersa*). When tightly massed in the perinuclear region of a photosensory cell as normally seen (Figure 1), or when compressed by centrifugation of an eye (REED & EAKIN, 1976), the vesicles become oval and packed into a paracrystalline array. Normally, large granules (35 nm or more) and a few mitochondria are dispersed among the vesicles. All other cytoplasmic structures appear to be shunted aside as the vesicles are produced and accumulated. The granules

¹ Dedicated to Dr. Rudolph Stohler, Founding Editor of *The Veliger*.

are considered beta particles of glycogen because of their size and digestibility by alpha amylase (EAKIN & BRANDENBURGER, 1967b). These features characterize only type I sensory cells of *H. aspersa*, not type II cells (BRANDENBURGER & EAKIN, 1974; BRANDENBURGER, 1975).

Origin and Fate

We believe that photic vesicles arise in the cell bodies of type I sensory cells by abscission of the ends of the Golgi cisternae—a conclusion based upon electron micrographs (EAKIN & BRANDENBURGER, 1967a) and upon osmium staining (EAKIN & BRANDENBURGER, 1970 see below). KATAOKA (1975), however, ascribes their origin to agranular endoplasmic reticulum (ER) in the slug *Limax flavus* because vesicles were sometimes observed in continuity with the smooth ER and because the vesicles, ER, and the convex side (but not the concave side) of the Golgi apparatus were stained with osmium tetroxide (see symposium discussion in EAKIN & BRANDENBURGER, 1976). In our experience, osmium staining revealed fine granular deposits in the smooth ER, whereas the Golgi apparatus and photic vesicles were intensely blackened. This observation accords with the prevailing dogma on the secretory activity of these organelles. After formation, photic vesicles appear to accumulate in large masses, described above. Then the vesicles move distally within the sensory cells. The movement is probably caused by pulsations of the eye resulting from contractions of unique musculo-secretory cells in the optic capsule (EAKIN & BRANDENBURGER, 1972; MORTENSEN & EAKIN, 1974). The final fate of the vesicles appears from our electron micrographs (EAKIN & BRANDENBURGER, 1982) and from osmium staining (Figure 2, EAKIN & BRANDENBURGER, 1970) to be fusion with smooth ER lying beneath the photoreceptor microvilli.

Transporters of Photopigment

One of the functions of photic vesicles is transport of photopigment along the pathway just described from perinuclear ER to the microvilli (EAKIN, 1972). Experiments, now presented in approximate chronological order, attest to this function.

Autoradiography: Snails (*Helix aspersa*) injected pericardially with tritiated vitamin A exhibited sequential uptake by Golgi, photic vesicles, and microvilli (EAKIN &



BRANDENBURGER, 1968; BRANDENBURGER & EAKIN, 1970). Although not definitively shown, it was assumed at that time that a snail's eye contains a photopigment whose chromophore is a vitamin A derivative. GILARY & WOLBARSH (1967) had found by electrical recording that the eye of another snail, *Otala lactea*, exhibited a spectral sensitivity in close agreement with the absorption curve of rhodopsin. Moreover, the Osaka workers, cited earlier, had discovered that green light (525 nm) was absorbed by the microvilli of a land snail (KOSHIDA *et al.*, 1963). Ultimately, OZAKI *et al.* (1986) demonstrated that the vesicles contain the photopigment retinochrome (discussed below). In our papers on the autoradiography of the eye of *H. aspersa*, mentioned above, we postulated that the aggregations of vesicles store a photopigment.

Osmium staining: In contrast to osmium fixation, staining with osmium tetroxide involves impregnation of specimens with the metal by 1–3 days of immersion in unbuffered 2% aqueous solution of OsO₄ at 40° C. When this procedure was applied to the eyes of *Helix aspersa*, the Golgi cisternae and photic vesicles were intensely stained (EAKIN & BRANDENBURGER, 1970). The so-called osmium black is believed to result from OsO₄ reduction by an unsaturated compound such as a vitamin A-containing photopigment. The rod disks of an amphibian (*Xenopus laevis*) were similarly stained. The distal ER beneath the microvilli and the bases of villi in the eyes of *H. aspersa* also showed intense osmiophilia (Figure 2), but distally the villi exhibited only a dusting of osmium black.

Light- and dark-adaptation: The detectable effects of light and darkness on photic vesicles suggest that the vesicles are transporters of photopigment (EAKIN & BRANDENBURGER, 1974). Snails (*Helix aspersa*) kept in total darkness for one to four months exhibited a break-up of the aggregations of vesicles, a marked increase in the production of lysosomes that incorporated the vesicles, and a deposition of multilayered membranous capsules and partitions around and within the masses of vesicles. These features indicated cessation in formation of photic vesicles and a removal of those in storage—a picture not observed in normal (wild) or light-adapted (control) snails. Moreover, in the latter an accumulation of vesicles distally near the microvilli was discernible (EAKIN & BRANDENBURGER, 1967c). The microvilli—the other principal organelles involved in photoreception—also showed ultrastructural changes in response to the presence or absence of light.

Explanation of Figures 1 and 2

Figure 1. Electron micrograph of photic vesicles massed near the nucleus of a type I photoreceptor cell in an eye of the snail *Helix aspersa*. G, glycogen granules. ×50,000. *Inset:* Some photic vesicles at higher magnification. ×100,000. Figure 2. Osmium staining of distal part of a type I photosensory cell in an eye of *Helix aspersa*. Note dense deposits of osmium black in photic vesicles (PV), endoplasmic reticulum (ER), and bases of some microvilli (MV). ×20,000. From EAKIN & BRANDENBURGER, 1970, with permission of *The Journal of Ultrastructure Research*.






Figure 3

Diagram of recycling of photopigment in eye of *Helix aspersa*. See text for explanation.

In another study, of a light-tolerant slug (*Ariolimax californicus*) and a nocturnal one (*Limax maximus*), we found a difference in the distribution of photic vesicles (EAKIN & BRANDENBURGER, 1975a). In the nocturnal species the vesicles were primarily in the perinuclear regions of the sensory cells, whereas in the diurnal species many vesicles occurred in the distal halves of the photoreceptor cells, as well as in masses in the cell bodies. All these observations support the hypothesis that light stimulates production, migration, and utilization of photic vesicles.

Autofluorescence: Knowing that vitamin A fluoresces (POPPER, 1944) and that the eyes of *Helix aspersa* contain this vitamin (EAKIN *et al.*, 1974), we examined sections of eyes of that snail by a fluorescence microscope to determine where the vitamin A was situated (EAKIN & BRANDENBURGER, 1978). The microvilli and the masses of photic vesicles immediately fluoresced, emitting a brilliant orange color that faded within 20–30 sec. Sections exposed to white light prior to examination under ultraviolet radiation did not fluoresce. As a control, sections of eyes of an amphibian (*Xenopus laevis*) were similarly treated and studied. The outer segments of the rods and cones fluoresced promptly, but the bright silvery color faded in 20–30 sec. Sections first exposed to white light did not fluoresce. Conclusion:

the photic vesicles may possess a vitamin A-containing compound such as a photopigment. In 1986 OZAKI *et al.* conducted a fluorescence microscopy study of the eye of a marine conch, *Conomurex luhuanus*, and observed autofluorescence of both microvilli and photic vesicles. There was a difference, however, in the light emitted by the two types of organelles.

Freeze-fracture: Carbon-platinum replicas of the surfaces of fractured, frozen eyes of *Helix aspersa* were studied by electron microscopy (BRANDENBURGER *et al.*, 1976) to obtain information on size and amount of intramembranous particles of photic vesicles and of photosensory microvilli. In light-adapted specimens (versus dark-adapted ones), 70 Å particles on the internal or protoplasmic (P) leaflet of the microvillar membranes were numerous. The particles, in the size range of rhodopsin, were interpreted as photopigment. Most photic vesicles cross-fractured. The vesicular membranes were usually not separated into P and E (external) leaflets, probably because of the small size of the vesicles. In favorable instances, however, in which the vesicular membrane was split, we found very few particles attached to either P or E surfaces. From this investigation I draw the conclusion that the putative photopigment, now known to be retinochrome (see below), is not carried in the vesicular membrane but in solution or colloidal sus-